

J. T. S.

Vol. 11 (2017), pp. 89-96

<https://doi.org/10.56424/jts.v11i01.10581>

On Concircularly ϕ –Recurrent Lorentzian α –Sasakian Manifold with Semi-Symmetric Non-Metric Connection

Kripasindhu Prasad and Sahadat Ali

Department of Mathematics,
Thakur Ram Multiple Campus, Birgunj Tribhuvan University, Nepal
Department of Mathematics, SRMGPC, Lucknow-227105, India
e-mail:kripasindhuchaudhary@gmail.com, sdalilko@rediffmail.com
(Received: January 02, 2017)

Abstract

The present study deals with concircular recurrent Lorentzian α –Sasakian manifolds with semi symmetric non-metric connection and discuss some interesting results.

1. Introduction

The notion of local symmetry of a Riemannian manifold has been weakened by many authors in several ways to a different extent. As a weaker version of local symmetry, Takahashi [12] introduced the notion of local ϕ –symmetry on a Sasakian manifold. Generalizing the notion of ϕ –symmetry, De, U. C. [6] introduced the notion of ϕ –recurrent Sasakian manifold.

Fridmann and Schouten introduced the idea of semi symmetric linear connection on a differentiable manifold. Hayden [1] introduced the idea of metric connection with torsion on Riemannian manifold. Further, Yano [14], Golab [8] defined and studied semi symmetric and quarter symmetric connection with affine connection, further many authors like De, U. C. [5], Sharfuddin and Hussain [2], Rastogi, Mishra and Pandey, Babewadi and many other studies the various properties of semi-symmetric connection.

In this paper we studied Concircular ϕ –recurrent Lorentzian α –Sasakian manifold with semi-symmetric non-metric connection and proved that a ϕ –recurrent Lorentzian α –Sasakian manifold with semi-symmetric non-metric connection is η –Einstein manifold. Further we have shown that in ϕ –recurrent Lorentzian α –Sasakian manifold with semi-symmetric non-metric connection the characteristic vector ξ and vector field ρ associated to the 1-form A are

codirectional. Finally, we proved that concircular ϕ -recurrent Lorentzian α -Sasakian manifold with semi-symmetric non-metric connection is η -Einstein manifold.

2. Preliminaries

A differentiable manifold M of dimension n is called a Lorentzian α -Sasakian manifold if it admits a tensor field ϕ of type $(1, 1)$, a contravariant vector field ξ , a covariant vector field η and Lorentzian metric g which satisfy

$$\phi^2 = 1 + \eta \otimes \xi, \quad (2.1)$$

$$\eta(\xi) = -1, \quad (2.2)$$

$$g(\phi X, \phi Y) = g(X, Y) + \eta(X)\eta(Y), \quad (2.3)$$

$$g(X, \xi) = \eta(X), \quad (2.4)$$

$$\phi\xi = 0, \quad \eta(\phi X) = 0, \quad (2.5)$$

$$(D_X \phi)Y = \alpha g(X, Y)\xi - \alpha \eta(Y)X, \quad (2.6)$$

for all $X, Y \in Tm$ [2, 3, 13].

Also a Lorentzian α -Sasakian manifold m satisfies

$$(D_X \xi)Y = \alpha \phi X, \quad (2.7)$$

$$(D_X \eta)Y = -\alpha g(\phi X, Y) \quad (2.8)$$

where D denotes the operator of covariant differentiation with respect to Lorentzian metric g . Also on a Lorentzian α -Sasakian manifold the following relation hold [2, 3, 13]

$$R(X, Y)\xi = \alpha^2(\eta(Y)X - \eta(X)Y), \quad (2.9)$$

$$R(\xi, X)Y = \alpha^2(g(X, Y)\xi - \eta(Y)X), \quad (2.10)$$

$$R(\xi, X)\xi = \alpha^2(\eta(X)\xi + X), \quad (2.11)$$

$$S(X, \xi) = (n - 1)\alpha^2\eta(X), \quad (2.12)$$

$$\eta(R(X, Y)Z) = \alpha^2(g(Y, Z)\eta(X) - g(X, Z)\eta(Y)), \quad (2.13)$$

$$g(R(\xi, X)Y, \xi) = -\alpha^2[g(X, Y) + \eta(X)\eta(Y)]. \quad (2.14)$$

For any vector field X, Y, Z where S is the Ricci curvature and Q is the Ricci operator given by

$$S(X, Y) = g(\phi X, Y).$$

A Lorentzian α -Sasakian manifold is said to be η -Einstein manifold if its Ricci tensor S takes the form

$$S(X, Y) = ag(X, Y) + b\eta(X)\eta(Y)$$

for arbitray vector X, Y where a and b are function on M . If $b = 0$, the η -Einstein manifold becomes Einstein manifold. [3,9] have proved that if Lorentzian α -Sasakian manifold M is η -Einstein manifold then $a+b = -\alpha^2(n-1)$.

Definition (2.1). A Lorentzian α -Sasakian manifold is said to be locally ϕ -symmetric if

$$\phi^2((D_W R)(X, Y)Z) = 0. \quad (2.15)$$

Definition (2.2). A Lorentzian α -Sasakian manifold is said to be recurrent if there exists a non zero 1-form A such that

$$\phi^2((D_W R)(X, Y)Z) = A(W)R(X, Y)Z, \quad (2.16)$$

where $A(W)$ is defined by $A(W) = g(W, \rho)$ and ρ is a vector field associated with the 1-form.

Definition (2.3). A Lorentzian α -Sasakian manifold is said to be concircularly recurrent if there exists a non zero 1-form A such that

$$\phi^2((D_W C)(X, Y)Z) = A(W)C(X, Y)Z, \quad (2.17)$$

where C is the concircular curvature tensor given by

$$C(X, Y)Z = R(X, Y)Z - \frac{r}{n(n-1)}[g(Y, Z)X - g(X, Z)Y], \quad (2.18)$$

where R is the Riemannian curvature tensor and r is the scalar curvature.

A linear connection D in an n -dimensional differentiable manifold is said to be semi-symmetric connection if its torsion tensor is of the form [1,5,8]

$$T(X, Y) = D_X Y - D_Y X - [X, Y] = \eta(Y)X - \eta(X)Y, \quad (2.19)$$

for all X, Y on TM . A semi-symmetric connection D is said to be semi-symmetric non-metric connection if it further satisfies $D_X g \neq 0$. [8]

3. Lorentzian α -Sasakian manifold with semi-symmetric non-metric connection

A semi-symmetric non-metric connection \bar{D} in Lorentzian α -Sasakian manifold can be defined by

$$\bar{D}_X Y = D_X Y + \eta(Y)X. \quad (3.1)$$

Also, we have

$$(\bar{D}_X g)(Y, Z) = -\eta(Y)g(X, Z) - \eta(Z)g(Y, X). \quad (3.2)$$

A connection given by (3.1) with (3.2) is called semi-symmetric non-metric connection in Lorentzian α -Sasakian manifold.

A relation between curvature tensor M of the manifold with semi-symmetric non-metric connection \bar{D} and Levi-Civita connection D is given by

$$\bar{R}(X, Y)Z = R(X, Y)Z - \alpha g(\phi X, Z)Y - \alpha g(\phi Y, Z)X, \quad (3.3)$$

where \bar{R} and R are the Riemannian curvature of the connection \bar{D} and D respectively.

From (3.3), we have

$$\bar{S}(Y, Z) = S(Y, Z) + \alpha(n-1)g(\phi Y, Z), \quad (3.4)$$

where \bar{S} and S are the Ricci tensor of the connection \bar{D} and D respectively.

Contracting (3.4), we get

$$\bar{r} = r, \quad (3.5)$$

where \bar{r} and r are the scalar curvatures of the connection \bar{D} and D respectively.

4. ϕ -Recurrent Lorentzian α -Sasakian manifold with semi-symmetric non-metric connection

Analogous to the definition (2.2) we define a Lorentzian α -Sasakian manifold is said to be ϕ -recurrent with respect to semi-symmetric non-metric connection if its curvature tensor \bar{R} satisfies the following condition

$$\phi^2((D_W \bar{R})(X, Y)Z) = A(W)\bar{R}(X, Y)Z. \quad (4.1)$$

Using (2.1) in (4.1), we get

$$(\bar{D}_W \bar{R})(X, Y)Z + \eta((\bar{D}_W \bar{R})(X, Y)Z)\xi = A(W)\bar{R}(X, Y)Z. \quad (4.2)$$

From which it follows that

$$g((\bar{D}_W \bar{R})(X, Y)Z, U) + \eta((\bar{D}_W \bar{R})(X, Y)Z)g(\xi, U) = A(W)g(\bar{R}(X, Y)Z, U). \quad (4.3)$$

Let $\{e_i\}$, $i = 1, 2, 3, \dots, n$ be an orthonormal basis of the tangent space at any point of the manifold. Then putting $X = U = \{e_i\}$ in (4.3) and taking summation over i , $1 \leq i \leq n$, we get

$$(\bar{D}_W \bar{S})(Y, Z) + \eta((\bar{D}_W \bar{R})(e_i, Y)Z)\eta(e_i) = A(W)\bar{S}(Y, Z). \quad (4.4)$$

Putting $Z = \xi$ in (4.4), the second term of (4.4) takes the form $g((\bar{D}_W \bar{R})(e_i, Y)\xi, \xi)$ which on simplification gives $g((\bar{D}_W \bar{R})(e_i, Y)\xi, \xi) = 0$.

Then from (4.4) we obtain

$$(\overline{D}_W \overline{S})(Y, \xi) = A(W) \overline{S}(Y, \xi). \quad (4.5)$$

Now, we know that

$$(\overline{D}_W \overline{S})(Y, \xi) = \overline{D}_W \overline{S}(Y, \xi) - \overline{S}(\overline{D}_W Y, \xi) - \overline{S}(Y, \overline{D}_W \xi). \quad (4.6)$$

Using (2.7), (2.8), (2.12), (3.4) in (4.6), we get

$$\begin{aligned} (\overline{D}_W \overline{S})(Y, \xi) = & \alpha S(Y, \phi W) + S(Y, W) - \alpha(\alpha + 1)(n - 1)g(Y, \phi W) \\ & - \alpha^2(n - 1)g(Y, W) + \alpha^2(n - 1)g(\phi Y, \phi W). \end{aligned} \quad (4.7)$$

In view of (4.5) and (4.7), we get

$$\begin{aligned} \alpha S(Y, \phi W) + S(Y, W) - \alpha(\alpha + 1)(n - 1)g(Y, \phi W) - \alpha^2(n - 1)g(Y, W) \\ + \alpha^2(n - 1)g(\phi Y, \phi W) = \alpha^2(n - 1)A(W)\eta(Y). \end{aligned}$$

Replacing $Y = \phi Y$ in above equation, we get

$$\begin{aligned} \alpha S(\phi Y, \phi W) + S(\phi Y, W) - \alpha(\alpha + 1)(n - 1)g(\phi Y, \phi W) \\ - \alpha^2(n - 1)g(\phi Y, W) + \alpha^2(n - 1)g(Y, \phi W) = 0. \end{aligned} \quad (4.8)$$

Interchanging Y and W in (4.8), we get

$$\begin{aligned} \alpha S(\phi W, \phi Y) + S(\phi W, Y) - \alpha(\alpha + 1)(n - 1)g(\phi W, \phi Y) \\ - \alpha^2(n - 1)g(\phi W, Y) + \alpha^2(n - 1)g(W, \phi Y) = 0. \end{aligned} \quad (4.9)$$

Adding (4.8) and (4.9) and simplifying, we get

$$S(\phi Y, \phi W) = (\alpha^2 + 1)(n - 1)g(\phi Y, \phi W).$$

Using (2.3) and (2.15), we get

$$S(Y, W) = (\alpha^2 + 1)(n - 1)g(Y, W) + (n - 1)\eta(Y)\eta(W).$$

This leads to the following theorem:

Theorem (4.1). A ϕ -recurrent Lorentzian α -Sasakian manifold with semi-symmetric non-metric connection in η -Einestien manifold.

Again (4.2), we have

$$(\overline{D}_W \overline{R})(X, Y)Z = -\eta((\overline{D}_W \overline{R})(X, Y)Z)\xi + A(W)\overline{R}(X, Y)Z. \quad (4.10)$$

From (2.13), (3.3) and using Bainchi identity, we get

$$A(W)\eta(\overline{R}(X, Y)Z) + A(X)\eta(\overline{R}(Y, W)Z) + A(Y)\eta(\overline{R}(W, X)Z) = 0. \quad (4.11)$$

From (2.13), (3.3) in (4.11), we get

$$A(W)\alpha^2[g(Y, Z)\eta(X) - g(X, Z)\eta(Y)] + A(X)\alpha^2[g(Z, W)\eta(Y)$$

$$\begin{aligned}
& -g(Y, Z)\eta(W)] + A(W)\alpha^2[g(X, W)\eta(Z) - g(Z, W)\eta(X)] \\
& + \alpha[g(\phi Y, Z)\eta(X) - g(\phi X, Z)\eta(Y) + g(\phi W, Z)\eta(Y) \\
& - g(\phi Y, Z)\eta(W) + g(\phi X, Z)\eta(W) - g(\phi W, Z)\eta(X)] = 0.
\end{aligned} \tag{4.12}$$

Putting $Y = Z = e_i$ in (4.12) and taking summation over i , $1 \leq i \leq n$, we get

$$A(W)\eta(X) = A(X)\eta(W) \tag{4.13}$$

for all vector fields W . Replacing X by ξ in (4.13), we get

$$A(W) = -\eta(\rho)\eta(W), \tag{4.14}$$

for any vector field W , where $A(\xi) = g(\xi, \rho) = \eta(\rho)$, ρ being vector field associated to the 1-form A that is $g(X, \rho) = A(X)$.

From (4.13) and (4.14), we state the following:

Theorem (4.2). In a ϕ -recurrent Lorentzian α -Sasakian manifold with semi-symmetric non-metric connection the characteristic vector ξ and vector field ρ associated to the 1-form A are codirectional and 1-form A is given by (4.14).

5. Concircular ϕ -Recurrent Lorentzian α -Sasakian manifold with Semi-symmetric Non-metric Connection

A analogous to the definition (2.3), we define a Lorentzian α -Sasakian manifold is said to be concircular ϕ -recurrent with respect to semi-symmetric non-metric connection if

$$\phi^2((D_W \bar{C})(X, Y)Z) = A(W)\bar{C}(X, Y)Z, \tag{5.1}$$

where \bar{C} is the concircular curvature tensor with respect to semi-symmetric non-metric connection given by

$$\bar{C}(X, Y)Z = \bar{R}(X, Y)Z - \frac{\bar{r}}{n(n-1)}[g(Y, Z)X - g(X, Z)Y]. \tag{5.2}$$

Using (2.1) in (5.1), we have

$$(\bar{D}_W \bar{C})(X, Y)Z + \eta((\bar{D}_W \bar{C})(X, Y)Z)\xi = A(W)\bar{C}(X, Y)Z. \tag{5.3}$$

From which it follows that

$$g((\bar{D}_W \bar{C})(X, Y)Z, U) + \eta((\bar{D}_W \bar{C})(X, Y)Z)g(\xi, U) = A(W)g(\bar{C}(X, Y)Z, U). \tag{5.4}$$

Let $\{e_i\}$, $i = 1, 2, 3, \dots, n$ be an orthonormal basis of the tangent space at any point of the manifold. Then putting $X = U = \{e_i\}$ in (4.3) and taking summation over i , $1 \leq i \leq n$, we get

$$(\overline{D}_W \overline{S})(Y, Z) = \frac{\overline{dr}(w)}{n} g(Y, Z) - \frac{\overline{dr}(w)}{n(n-1)} [g(Y, Z) - \eta(Y)\eta(Z)] \\ + A(W) \left[\overline{S}(Y, Z) - \frac{\bar{r}}{n} g(Y, Z) \right].$$

Putting $Z = \xi$ and using (2.2) and (2.4) in above, we have

$$(\overline{D}_W \overline{S})(Y, \xi) = \frac{\overline{dr}(w)}{n} \eta(Y) - \frac{\overline{dr}(w)}{n(n-1)} 2\eta(Y) + A(W) \left[\overline{S}(Y, \xi) - \frac{\bar{r}}{n} \eta(Y) \right]. \quad (5.5)$$

Now, we know that

$$(\overline{D}_W \overline{S})(Y, \xi) = \overline{D}_W \overline{S}(Y, \xi) - \overline{S}(\overline{D}_W Y, \xi) - \overline{S}(Y, \overline{D}_W \xi). \quad (5.6)$$

Using (2.7) (2.8), (2.12), (3.4) in (5.6), we get

$$(\overline{D}_W \overline{S})(Y, \xi) = \alpha S(Y, \phi W) + S(Y, W) - \alpha(\alpha+1)(n-1)g(Y, \phi W) \\ - \alpha^2(n-1)g(Y, W) + \alpha^2(n-1)g(\phi Y, \phi W). \quad (5.7)$$

In view of (5.5) and (5.7), we get

$$\alpha S(Y, \phi W) + S(Y, W) - \alpha(\alpha+1)(n-1)g(Y, \phi W) - \alpha^2(n-1)g(Y, W) \\ + \alpha^2(n-1)g(\phi Y, \phi W) = \frac{\overline{dr}(w)}{n} \eta(Y) - \frac{\overline{dr}(w)}{n(n-1)} 2\eta(Y) \\ + A(W) \left[\overline{S}(Y, \xi) - \frac{\bar{r}}{n} \eta(Y) \right].$$

Replacing $Y = \phi Y$ in above equation, we get

$$\alpha S(\phi Y, \phi W) + S(\phi Y, W) - \alpha(\alpha+1)(n-1)g(\phi Y, \phi W) \\ - \alpha^2(n-1)g(\phi Y, W) + \alpha^2(n-1)g(Y, \phi W) = 0. \quad (5.8)$$

Interchanging Y and W in (5.8), we get

$$\alpha S(\phi W, \phi Y) + S(\phi W, Y) - \alpha(\alpha+1)(n-1)g(\phi W, \phi Y) \\ - \alpha^2(n-1)g(\phi W, Y) + \alpha^2(n-1)g(W, \phi Y) = 0. \quad (5.9)$$

Adding (5.8) and (5.9) and simplifying we get

$$S(\phi Y, \phi W) = (\alpha^2 + 1)(n-1)g(\phi Y, \phi W).$$

Using (2.3) and (2.15), we get

$$S(Y, W) = (\alpha^2 + 1)(n-1)g(Y, W) + (n-1)\eta(Y)\eta(W).$$

This leads to the following theorem:

Theorem (5.1). A concircular ϕ -recurrent Lorentzian α -Sasakian manifold with semi-symmetric non-metric connection is η -Einestien manifold.

REFERENCES

- [1] Sharfuddin, A. and Husain, S. I. : Semi symmetric metric connection in almost contact manifold, Tensor, 30(2)(1976), 133-139.
- [2] Bhattacharya, A. and Tarafdar, M. : On Lorentzian para sasakian manifolds, Steps in differential geometry, In: Proc. of the Coll. On differential Geo., Debreon (2000).
- [3] Yildiz, A. and Murathan, C. : On Lorentzian α -Sasakian manifolds, Kyungpook Math. J., 45 (2005), 95-103.
- [4] De, U. C. : On a type of semi symmetric metric connection on Riemannian manifold, Indian J. Pure Appl. Math., 21(4)(1990), 334-338.
- [5] De, U. C., Shaikh, A. A. and Biswas, S. : On ϕ -Recurrent Sasakian manifolds, Novi Sad J. Math, 33(2) (2003), 43-48.
- [6] De, U. C. : On ϕ -symmetric Kenmotsu manifold. Int. Electronic J. Geometry, 1(1)(2008), 33-38.
- [7] Golab, S. : On semi symmetric and quarter symmetric linear connections, Tensor, 29(3) (1975), 249-254.
- [8] Kenmotsu, K. : A class of almost contact Riemannian manifolds, Tohoku Math. J., 24 (1972), 93-103.
- [9] Sato, I. : On structure similar to the almost contact structure, Tensor, 30 (1976), 219-224.
- [10] Srivastava, S. K. and Prakash, A. : On concircularly ϕ -recurrent Sasakian manifolds, Globale J. of Pure and Applied Mathematics, 9(2)(2013), 215-220.
- [11] Takahashi, T. : Sasakian ϕ -symmetric spaces, Tohoku Math. J., 29 (1977), 91-113.
- [12] Taleshin and Asghari, N. : On Lorentzian α -Sasakian manifolds, The J. of Mathematics and Computer Science, 14(3)(2012), 295-300.
- [13] Yano, K. : On semi-symmetric metric connection, Reveu Roumaine de Mathematiques Pures et., 15 (1970), 1579-1586.
- [14] Yano, K. : Concircular geometry-1, concircular transformation, Proceedings of the Japan Academy, 16 (1940), 195-200.